



## Cytokeratin 7 (RCK105)

**CATALOG NUMBER:** MUB0315P

**CLONE:** RCK105

**SPECIES / ISOTYPE:** mouse IgG1

**PRODUCT FORM:** purified monoclonal antibody

### BACKGROUND

Cytokeratins are a subfamily of intermediate filament proteins and are characterized by a remarkable biochemical diversity, represented in human epithelial tissues by at least 20 different polypeptides. They range in molecular weight between 40 kDa and 68 kDa and isoelectric pH between 4.9 – 7.8. The individual human cytokeratins are numbered 1 to 20.

The various epithelia in the human body usually express cytokeratins which are not only characteristic of the type of epithelium, but also related to the degree of maturation or differentiation within an epithelium.

Cytokeratin subtype expression patterns are used to an increasing extent in the distinction of different types of epithelial malignancies. The cytokeratin antibodies are not only of assistance in the differential diagnosis of tumors using immunohistochemistry on tissue sections, but are also a useful tool in cytopathology and flow cytometric assays.

### SOURCE

RCK105 is a mouse monoclonal IgG1 antibody derived by fusion of SP2/0-Ag14 mouse myeloma cells with spleen cells from a BALB/c mouse immunized with cytokeratins from the human bladder carcinoma cell line T24.

### PRODUCT

Each vial contains 100 µl 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.

### SPECIFICITY

RCK105 reacts exclusively with cytokeratin 7 which is present in a subgroup of glandular epithelia and their tumors, as well as transitional epithelium and transitional carcinoma.

RCK105 is suitable for immunoblotting, immunocytochemistry, immunohistochemistry on frozen sections and flow cytometry. Optimal antibody dilution should be determined by titration; recommended range is 1:100 – 1:200 for flow cytometry, and for immunohistochemistry with avidin-biotinylated horseradish peroxidase complex (ABC) as detection reagent, and 1:100 – 1:1000 for immunoblotting applications.

### SPECIES REACTIVITY

Human, mouse, rat, hamster, canine, swine and feline.

### STORAGE

Store at 4°C, or in small aliquots at –20°C.

### REFERENCES

1. Ramaekers, F., Huysmans, A., Schaart, G., Moesker, O., and Vooijs, P. (1987). Tissue distribution of keratin 7 as monitored by a monoclonal antibody, *Exp Cell Res* 170, 235-49.
2. Smedts, F., Ramaekers, F., Robben, H., Pruszczynski, M., van Muijen, G., Lane, B., Leigh, I., and Vooijs, P. (1990). Changing patterns of keratin expression during progression of cervical intraepithelial neoplasia, *Am J Pathol* 136, 657-68.
3. Schaafsma, H. E., Ramaekers, F. C., van Muijen, G. N., Lane, E. B., Leigh, I. M., Robben, H., Huijsmans, A., Ooms, E. C., and Ruiten, D. J. (1990). Distribution of cytokeratin polypeptides in human transitional cell carcinomas, with special emphasis on changing expression patterns during tumor progression, *Am J Pathol* 136, 329-43.
4. Ramaekers, F., van Niekerk, C., Poels, L., Schaafsma, E., Huijsmans, A., Robben, H., Schaart, G., and Vooijs, P. (1990). Use of monoclonal antibodies to keratin 7 in the differential diagnosis of adenocarcinomas, *Am J Pathol* 136, 641-55.
5. Bauwens, L. J., De Groot, J. C., Ramaekers, F. C., Veldman, J. E., and Huizing, E. H. (1992). Expression of intermediate filament proteins in the adult human vestibular labyrinth, *Ann Otol Rhinol Laryngol* 101, 479-86.
6. Smedts, F., Ramaekers, F., Troyanovsky, S., Pruszczynski, M., Link, M., Lane, B., Leigh, I., Schijf, C., and Vooijs, P. (1992). Keratin expression in cervical cancer, *Am J Pathol* 141, 497-511.
7. Ivanyi, D., Minke, J. M., Hageman, C., Groeneveld, E., and van Doornwaard, G. (1992). Patterns of expression of feline cytokeratins in healthy epithelia and mammary carcinoma cells, *Am J Vet Res* 53, 304-14.

© 2009 MUBio Products B.V.

Datasheet version: MUB\_0315P\_090825

### WARNING and CAUTION

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals.

This product contains sodium azide. To prevent formation of toxic vapors, do not mix with strong acidic solutions. To prevent formation of potentially explosive metallic azides in metal plumbing, always wash into drain with copious quantities of water.

This datasheet is as accurate as reasonably achievable, but MUBio Products BV accepts no liability for any inaccuracies or omissions in this information.