



Cytokeratin 10 (RKSE60)

CATALOG NUMBER: MUB0319P
CLONE: RKSE60
SPECIES / ISOTYPE: mouse IgG1
PRODUCT FORM: purified monoclonal antibody

BACKGROUND

Cytokeratins are a subfamily of intermediate filament proteins and are characterized by a remarkable biochemical diversity, represented in human epithelial tissues by at least 20 different polypeptides. They range in molecular weight between 40 kDa and 68 kDa and isoelectric pH between 4.9 – 7.8. The individual human cytokeratins are numbered 1 to 20.

The various epithelia in the human body usually express cytokeratins which are not only characteristic of the type of epithelium, but also related to the degree of maturation or differentiation within an epithelium.

Cytokeratin subtype expression patterns are used to an increasing extent in the distinction of different types of epithelial malignancies. The cytokeratin antibodies are not only of assistance in the differential diagnosis of tumors using immunohistochemistry on tissue sections, but are also a useful tool in cytopathology and flow cytometric assays.

SOURCE

RKSE60 is a mouse monoclonal IgG1 antibody derived by fusion of SP2/0 mouse myeloma cells with spleen cells from a mouse immunized with cytokeratins from the human epidermis.

PRODUCT

Each vial contains 100 µl 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.

SPECIFICITY

RKSE60 reacts exclusively with cytokeratin 10 which is present in keratinizing stratified epithelia and in differentiated areas of highly differentiated squamous cell carcinomas.

RKSE60 is suitable for immunoblotting, immunocytochemistry, immunohistochemistry on frozen tissues and flow cytometry. Optimal antibody dilution should be determined by titration; recommended range is 1:100 – 1:200 for flow cytometry, and for immunohistochemistry with avidin-biotinylated horseradish peroxidase complex

(ABC) as detection reagent, and 1:100 – 1:1000 for immunoblotting applications.

SPECIES REACTIVITY

Human, mouse, rat, canine.

STORAGE

Store at 4°C, or in small aliquots at –20°C.

REFERENCES

1. Ramaekers, F. C., Puts, J. J., Moesker, O., Kant, A., Huysmans, A., Haag, D., Jap, P. H., Herman, C. J., and Vooijs, G. P. (1983). Antibodies to intermediate filament proteins in the immunohistochemical identification of human tumours: an overview, *Histochem J* 15, 691-713.
2. Puts, J. J., Moesker, O., Kenemans, P., Vooijs, G. P., and Ramaekers, F. C. (1985). Expression of cytokeratins in early neoplastic epithelial lesions of the uterine cervix, *Int J Gynecol Pathol* 4, 300-13.
3. Broers, J. L., Carney, D. N., Klein Rot, M., Schaart, G., Lane, E. B., Vooijs, G. P., and Ramaekers, F. C. (1986). Intermediate filament proteins in classic and variant types of small cell lung carcinoma cell lines: a biochemical and immunochemical analysis using a panel of monoclonal and polyclonal antibodies, *J Cell Sci* 83, 37-60.
4. Ramaekers, F., Huysmans, A., Schaart, G., Moesker, O., and Vooijs, P. (1987). Tissue distribution of keratin 7 as monitored by a monoclonal antibody, *Exp Cell Res* 170, 235-49.
5. Van Muijen, G. N., Warnaar, S. O., and Ponc, M. (1987). Differentiation-related changes of cytokeratin expression in cultured keratinocytes and in fetal, newborn, and adult epidermis, *Exp Cell Res* 171, 331-45.
6. Ramaekers, F., van Niekerk, C., Poels, L., Schaafsma, E., Huijsmans, A., Robben, H., Schaart, G., and Vooijs, P. (1990). Use of monoclonal antibodies to keratin 7 in the differential diagnosis of adenocarcinomas, *Am J Pathol* 136, 641-55.
7. Schaafsma, H. E., Ramaekers, F. C., van Muijen, G. N., Lane, E. B., Leigh, I. M., Robben,

WARNING and CAUTION

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals.

This product contains sodium azide. To prevent formation of toxic vapors, do not mix with strong acidic solutions. To prevent formation of potentially explosive metallic azides in metal plumbing, always wash into drain with copious quantities of water.

This datasheet is as accurate as reasonably achievable, but MUBio Products BV accepts no liability for any inaccuracies or omissions in this information.



- H., Huijsmans, A., Ooms, E. C., and Ruiters, D. J. (1990). Distribution of cytokeratin polypeptides in human transitional cell carcinomas, with special emphasis on changing expression patterns during tumor progression, *Am J Pathol* 136, 329-43.
8. Smedts, F., Ramaekers, F., Robben, H., Pruszczynski, M., van Muijen, G., Lane, B., Leigh, I., and Vooijs, P. (1990). Changing patterns of keratin expression during progression of cervical intraepithelial neoplasia, *Am J Pathol* 136, 657-68.

© 2009 MUBio Products B.V.
Datasheet version: MUB_0319P_090604

WARNING and CAUTION

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals.
This product contains sodium azide. To prevent formation of toxic vapors, do not mix with strong acidic solutions. To prevent formation of potentially explosive metallic azides in metal plumbing, always wash into drain with copious quantities of water.